

# Profile distribution of soil organic carbon under different land use type in Sanjing Plain

Chi Guangyu, Chen Xin, Shi Yi

Key Laboratory of Terrestrial Ecological Process, Institute of Applied Ecology, Chinese Academy of Sciences, Email Chi Guangyu, [chiguangyu1018@126.com](mailto:chiguangyu1018@126.com), Chen Xin, [chenxin@iae.ac.cn](mailto:chenxin@iae.ac.cn), Shi Yi, [shiyi@iae.ac.cn](mailto:shiyi@iae.ac.cn)

## Abstract

To understand the dynamic changes of soil organic carbon (SOC) after different durations of cultivation, soil samples down to a depth of 120 cm were collected in layers from lowland and upland fields having been reclaimed for 5-25 years, with adjacent undisturbed wetland and forestland as the controls. The study of the vertical distribution of SOC and its relationship with soil pH showed that the SOC content in undisturbed wetland and cultivated lowland rice fields had a marked decrease from 0-10 cm to 40-60 cm and a smaller change downward, and a similar variation trend was observed in undisturbed forestland and cultivated soybean fields, only with the difference that the SOC content in 0-10 cm layer was much higher in forestland than in wetland, and lower in soybean fields than in rice fields. For undisturbed wetland, the SOC content in surface layer was decreased by 49.3% and 14.3% after being reclaimed for 10 and 25 years, and for undisturbed forestland, 81.9% and 68.3% of SOC in surface layer were lost after being reclaimed for 5 years and 18 years, respectively.

## Key Words

Soil organic carbon (SOC), pH, land use, Sanjing Plain

## Introduction

Soil organic carbon (SOC) is a main factor affecting soil quality and agriculture sustainability. Being a source and sink of plant nutrients, SOC plays an important role in terrestrial C cycle (Freixo *et al.*, 2002). Land use type has a deep effect on SOC storage, since it affects the amount and quality of litter input, litter decomposition rate, and stabilization of SOC. The SOC loss from irrational land use often leads to some negative impacts on both terrestrial and aquatic ecosystems, and on atmospheric environment (Reeder *et al.*, 1998; Bronson *et al.*, 2004). The Sanjiang Plain, one of the largest freshwater marshes in China, has been experienced intensive cultivation over past 50 years. About 3.8 Mha of its native marshland has been converted into cultivated land, resulting in a significant change in hydrological properties of the Plain (Liu and Ma, 2000). Many researches were made on the dynamics of methane emission due to this land use change (Ding *et al.*, 2004), but the effects of the land conversion on SOC remain largely unknown. With the cultivated lowland and upland and adjacent undisturbed wetland and forestland as test objects, this paper studied the dynamic changes of SOC under different land use type in Sanjiang Plain.

## Methods

### Sampling sites

The undisturbed wetland and forestland and cultivated lowland and upland in Sanjiang Honghe Farm were selected as test objects, with their sampling sites listed in Table 1.

Table 1. Description of sampling sites.

No.	Land use type	Reclamation history
Site 1	Wetland	Undisturbed
Site 2	Lowland rice field	Reclaimed for 10 years
Site 3	Lowland rice field	Reclaimed for 25 years
Site 4	Forestland	Undisturbed
Site 5	Upland soybean field	Reclaimed for 5 years
Site 6	Upland soybean field	Reclaimed for 18 years

### Soil sampling and analysis


Soil samples were taken from the depths of 0-10 cm, 10-20 cm, 20-40 cm, 40-60 cm, 60-90 cm and 90-120 cm, and duplicates were installed at each sampling site. Soil organic carbon content was determined by TOC

5000A autoanalyzer, and soil pH was measured in 1:2.5 soil:water suspension by using Elico Digital EC meter. Statistic analyses were made with SPSS 10.0.

## Results

### Soil organic carbon

The SOC content in undisturbed wetland and cultivated lowland rice fields had a marked decrease from 0-10 cm to 40-60 cm and a less change downward (Figure 1a). In 0-10 cm layer, there was a significant difference in SOC content ( $P < 0.01$ ), with the sequence of undisturbed wetland > lowland rice field reclaimed for 25 years > lowland rice field reclaimed for 10 years. Compared with that in undisturbed wetland, the SOC content in the 0-10 cm layer in the lowland rice fields having been reclaimed for 10 and 25 years was decreased by 49.3% and 14.3%, respectively.



**Figure 1. Vertical distribution of SOC under different land use types in Sanjiang Plain.**


A similar distribution pattern of SOC was observed in undisturbed forestland and cultivated soybean fields (Figure 1b), only with the difference that the SOC content in the 0-10 cm layer was much higher in forestland than in wetland, and lower in soybean fields than in lowland rice fields. Compared with that in undisturbed forestland, the SOC content in the 0-10 cm layer in the soybean fields having been reclaimed for 5 and 18 years was decreased by 81.9% and 68.3%, respectively.

The higher storage of SOC in surface layer was closely related with the accumulation of plant materials, while the differences in the dynamics of SOC in this layer should have close relations with the amount and quality of plant residues, as well as the environmental and soil conditions (Needelman *et al.*, 1999).

### Soil pH

Soil pH decreased with depth in undisturbed wetland, but had a uniform vertical distribution in cultivated lowland rice fields. It was higher throughout the profile in the rice field with a longer reclamation history than in that with a shorter one. In 0-10 cm layer, there was a significance difference in soil pH, with the sequence of undisturbed wetland > lowland rice field reclaimed for 25 years > lowland rice field reclaimed for 10 years (Figure 2a). On the contrary, the soil pH in undisturbed forestland and the soybean field having been reclaimed for 18 years was increased with depth, and in 0-10 cm layer, soil pH was decreased in the sequence of soybean field reclaimed for 5 years > soybean field reclaimed for 18 years > undisturbed forestland (Figure 2b).

The different distribution patterns of soil pH suggested that reclamation had different effects on soil acidity of wetland and forestland, especially that in surface layer, which should have definite effects on SOC storage. Regression analysis revealed that there was a significant negative correlation between soil pH and SOC in undisturbed forestland and the soybean field having been reclaimed for 18 years, indicating that the higher soil pH after reclamation led to a decreased SOC storage, probably due to the enhanced mineralization of SOC by soil microbes (Motavalli *et al.*, 1995).



**Figure 2. Vertical distribution of soil pH under different land use type in Sanjiang Plain.**

### Conclusion

A similar vertical distribution pattern of SOC, *i.e.*, decreased markedly from 0-10 cm to 40-60 cm and less changed downward, was observed in undisturbed wetland and forestland and in their reclaimed fields. The only difference was that the SOC content in 0-10 cm layer was much higher in forestland than in wetland, and lower in soybean fields than in rice fields. Reclamation made a great loss of SOC in surface layer, with a loss rate of 49.3% and 14.3% in wetland after reclaimed for 10 and 25 years, and 81.9% and 68.3% in forestland after reclaimed for 5 years and 18 years, respectively.

Land use type had a significant effect on soil pH. In surface layer, soil pH was decreased in the sequences of undisturbed wetland > lowland rice field reclaimed for 25 years > lowland rice field reclaimed for 10 years, and soybean field reclaimed for 5 years > soybean field reclaimed for 18 years > undisturbed forestland. The variations of surface soil pH under reclamation could be one of the factors inducing the SOC loss.

### References

- Bronson K, Zobeck T, Chua TT, Acosta-Martinez V, van Pelt RS, Booker JD (2004) Carbon and nitrogen pools of southern high plains cropland and grassland soils. *Soil Sci. Soc. Am. J.* **68**, 1695-1704.
- Ding WX, Cai ZC, Tsuruta H (2004) Cultivation, nitrogen fertilization and set-aside effects on methane uptake in a drained marsh soil in Northeast China. *Glob. Change Biol.* **10**, 1801-1809.
- Freixo AA, Machado PL, Santos HP, Silva CA, Fadigas FS (2002) Soil organic carbon and fractions of a Rhodic Ferralsol under the influence of tillage and crop rotation systems in southern Brazil. *Soil Tillage Res.* **64**, 221-230.
- Hassink J, Whitmore AP, Kubat J (1997) Size and density fractionation of soil organic matter and the physical capacity of soils to protect organic matter. *Eur. J. Agron.* **7**, 189-199.
- Liu ZG, Ma XH (1997) Effect of reclamation on soil environment in Sanjiang Plain. *Pedosphere* **7**, 73-78.
- Marsh Research Department, Changchun Institute of Geography, Academia Sinica. Marsh of Sanjiang Plain (in Chinese). Science Press, Beijing. 1983: 208.
- Rimkens PFAM, van der PJ, Hassink J (1999) Soil organic matter dynamics after the conversion of arable land to pasture. *Biol. Fertil. Soils* **28**, 277-284.
- Reeder JD, Schumman GE, Bowman RA (1998) Soil C and N changes on conservation reserve program lands in the Central Great Plains. *Soil Tillage Res.* **47**, 339-349.
- Six J, Callewaert P, Lenders S, de Gryze S, Morris SJ, Grogovich EG, Paul EA, Paustian K (2002) Measuring and understanding carbon storage in afforested soils by physical fractionation. *Soil Sci. Soc. Am. J.* **66**, 1981-1987.
- Six J, Paustian K, Elliott ET, Combrink C (2000) Soil structure and organic matter: I. Distribution of aggregates-size classes and aggregate-associated carbon. *Soil Sci. Soc. Am. J.* **37**, 509-513.
- Shepherd TG, Saggarr S, Newman RH, Ross CW, Dando JL (2001) Tillage-induced changes to soil structure and organic carbon fractions in New Zealand soils. *Aust. J. Soil Res.* **39**, 465-489.